



The Relationship Between Postoperative Pain Intensity and Sleep Quality in Patients After Spinal Anesthesia

journal home page: <https://goicare.web.id/index.php/JNJ>

Shelina¹, Amin Susanto¹, Septian Mixrova Sebayang¹

¹Anesthesiology Nursing Study Program Applied Undergraduate Program,
Faculty of Health Harapan Bangsa University



CROSS-SECTIONAL STUDY

ARTICLE HISTORY

Received: September 16, 2024

Revised: November 1, 2024

Accepted: February 9, 2025

DOI: 10.61716/jnj.v3i1.97

*Corresponding author:

Shelina

Anesthesiology Nursing Study Program
Undergraduate Program Faculty of
Health Harapan Bangsa University,
Indonesia Jl. Raden Patah No.100,
Ledug, Kembaran, Purwokerto,
Banyumas, Central Java, Indonesia.

Email: shelinapark30@gmail.com



Abstract

Background: Postoperative pain in each patient will experience different pain from one another. Inadequate pain management can have a direct impact on postoperative pain, which is disruption of sleep quality. **Purpose:** The purpose of this research was to identify any correlation between the severity of postoperative pain and the quality of sleep experienced by patients after spinal anesthesia. **Methods:** This research employs a cross-sectional design based on quantitative correlation. Using a non-probability purposive sampling approach, 51 participants were surveyed after spinal anesthesia for this investigation. The data was collected in the inpatient room at Wijayakusuma Purwokerto Hospital using questionnaire sheets from the Pittsburgh Sleep Quality Index (PSQI) to assess sleep quality and NRS (Numeric Rating Scale) to quantify postoperative pain severity. The Spearman rank test was used to examine the data. **Results:** Forty-three people (78.4% of the total) reported minor postoperative discomfort, while forty-three people (84.3% of the total) reported satisfactory sleep quality. A correlation force of 0.756 and a p-value of 0.000 were determined using the Spearman rank test. **Conclusions:** Patients undergoing spinal anesthesia at Wijayakusuma Purwokerto Hospital reported significantly higher levels of postoperative discomfort and worse sleep quality.

Keywords: pain intensity; sleep quality; post-operative; spinal anesthesia

Introduction

Surgery or surgery is one of the invasive actions carried out by making an incision in a part of the body that is experiencing health problems, then taking corrective action which ends with closing the wound through the suturing process [1]. In its implementation, surgery requires anesthesia which aims to relieve surgical pain [2].

Anesthesia has 3 types, namely local anesthesia, regional anesthesia and general anesthesia. The anesthetic technique that is often used in lower extremity surgery is regional anesthesia with a spinal technique (Subarachnoid

Block). Spinal anesthesia in the process of which local anesthetic drugs are injected into the subarachnoid space between the lumbar vertebrae 2-3, lumbar 3-4 or lumbar 4-5 [3]. The typical duration of spinal anesthetic treatments is around 4 to 6 hours, although this might vary depending on a number of circumstances. After 4 to 6 hours the analgesic effect of spinal anesthesia drugs begins to decrease so that patients will feel pain in the surgical wound. Post-anesthesia patients will experience pain even though pharmacological therapy has been given, especially during the interval of analgesic administration [4].

The complicated response of the Central Nervous System (CNS) to tissue stress causes postoperative pain [5]. The pain can be mild, moderate, or severe; however, as the body heals, the intensity of the pain may decrease. Delayed rehabilitation and prolonged hospitalization can be caused by uncontrolled patient pain [6].

Pain perception varies greatly in each individual, this can occur because pain is subjective [7]. Discomfort is a uniquely personal experience that no one else can understand. Pain can fill a person's entire mind, regulate his activities and change the person's life [8]. Pain management can be done for post-surgical pain management with pharmacological and non-pharmacological therapies [9,10]. The implementation of analgesic administration is most often given to patients who experience severe pain [1]. Therefore, adequate post-surgical pain treatment is very important [11]. Inadequate pain management can have effects on patients, such as sleep disturbances, difficulty mobilizing, anxiety and aggression [1]. Disruption of sleep quality with improper rapid eye movement (REM) and non-REM (nonrapid eye movement) periods is a direct result of the pain that will be experienced following surgery [12].

Sleep disruption is characterized by a lack of rejuvenation upon awakening or difficulty going asleep, even while the total amount of time spent sleeping is met. Sleep disorders can cause decreased pain tolerance due to increased fatigue in the sympathetic central nervous system which leads to increased analgesic use which contributes to the incidence of sleep disorders [13].

According to a study conducted by Budiyo and Hamdiah (2022), a total of 12 participants (63.2%) reported excellent sleep quality while experiencing light pain, 17 participants (73.9%) reported moderate

pain, and 15 participants (93.8%) reported severe pain while experiencing poor sleep quality. While one respondent (5.3%) had poor sleep quality and eighteen (94.7%) had excellent sleep quality out of nineteen respondents with moderate pain levels in the study by [14,15]

Data from the past three months with an average of 106 patients having spinal anaesthetic is available, according to a pre-survey done at RST Wijayakusuma Purwokerto on November 21, 2023. Supported by the results of an interview with one of the patients after spinal anesthesia surgery after 24 hours that there were complaints of mild pain after surgery.

Based on the description above, there has not been much research on the phenomenon that occurs and the information that has been obtained, The researcher is keen in delving into this phenomenon via study at RST Wijayakusuma Purwokerto, specifically looking for a correlation between the severity of postoperative pain and the quality of sleep experienced by patients after spinal anesthesia.

Methods

This study employs a cross-sectional design based on quantitative correlation. After spinal anesthesia, 51 participants were selected using a non-probabilistic purposive selection method for this investigation. Patients undergoing spinal anesthesia at RST Wijayakusuma Purwokerto after surgery made up the study's population. On a monthly basis, during August and October 2023, RST Wijayakusuma Purwokerto averaged 106 patients undergoing spinal anesthesia for surgical procedures. Using the numerical Rating Scale (NRS) and the Pittsburgh Sleep Quality Index (PSQI), data was collected from June 28 to July 13, 2024, in the Antasena and Gayatri rooms at RST Wijayakusuma Purwokerto to assess the

severity of postoperative pain and sleep quality, respectively. Applying the spearman rank test to the data. Approval number B.L.PPM-UHB/557/06/2024 it indicates that this study has been greenlit by the University of Harrogate's Research Ethics Committee.

Findings

Table 1: Characteristics of Respondents following Spinal Anesthesia at RST Wijayakusuma Purwokerto: Frequency Distribution

Characteristics	Frequency (n=51)	Percentage
Age		
18-30 years	18	35,3
31-40 years	9	17,6
41-50 years	9	17,6
51-60 years	11	21,6
> 60 years	4	7,8
Gender		
Male	22	43,1
Female	29	56,9

Table 1 shows the demographics of the 51 respondents, broken down by gender and age. Age characteristics show that at the age of 18-30 years as many as 18 respondents (35%), 31-40 years old as many as 9 respondents (17.6%), 41-50 years of age 9 respondents (17.6%), 51-60 years old as many as 11 respondents (21.6%), and at the age of >60 years as many as 4 respondents (7.8%). Whereas in gender characteristics in women as many as 29 respondents (56.9%) and in men as many as 22 respondents (43.2%).

Table 2. Distribution of Postoperative Pain Intensity in Patients at RST Wijayakusuma Purwokerto Following Spinal Anesthesia

Pain intensity	Frequency (n=51)	Percentage (100%)
No pain (0)	3	5,9
Mild pain (1-3)	40	78,4
Moderate pain (4-6)	8	15,7

Severe pain (7-10)	0	0
--------------------	---	---

According to Table 2, out of 51 respondents, 3 (or 5.9%) reported no pain at all, 40 (or 78.4%) reported mild pain, 8 (15.7%) reported moderate pain, and 0 (or 0%) reported severe pain after the operation.

Table 3. Data from patients undergoing spinal anesthesia at RST Wijayakusuma Purwokerto on the frequency distribution of sleep quality

Sleep Quality	Frequency (n=51)	Percentage (100%)
Good sleep quality	43	84,3
Poor sleep quality	8	15,7

According to Table 3, which displays the findings of the frequency distribution of sleep quality, 44 respondents (84.3%) had excellent sleep quality, whereas 8 respondents (15.7%) had poor sleep quality.

Table 4. Relationship between Postoperative Pain Intensity and Sleep Quality in Patients After Spinal Anesthesia at RST Wijayakusuma Purwokerto.

Pain Intensity	Sleep Quality				p-value	r
	Good n	%	Bad n	Total n %		
No pain (0)	3	5,9	0	0,0	3	5,9
Mild pain (1-3)	39	76,5	1	2,0	40	78,4
Moderate pain (4-6)	1	2,0	7	13,7	8	15,7
Severe pain	0	0	0	0	0	0

(7-10)

Total	4 3	84, 3	8	15, 7	51	10 0
-------	--------	----------	---	----------	----	---------

According to Table 4, the number of respondents who reported no discomfort while sleeping is 5.9%, and there was not a single respondent who reported poor sleep quality. With mild discomfort, 36 people (76.5%) reported poor sleep quality, while 1 person (2.0%) reported poor sleep quality. One respondent (2.0%) reported excellent sleep quality while suffering moderate discomfort, whereas seven (13.7%) reported poor sleep quality.

Discussions

According to table 1 the majority of patients in this research who had spinal anesthesia for surgery were between the ages of 18 and 30, comprising 18 respondents or 35.5% of the total. This is in line with Sugathot and Utami's research (2018) which says age is an important variable that affects pain, especially in children and the elderly. Adults will experience neurophysiological changes and may experience a decrease in sensory stimulus perception and an increase in pain threshold. The increasing age, the higher the pain reaction and response felt [16]. Based on the characteristics of gender in this study, the majority occurred in women as many as 29 respondents (56.9%). This is due to the fact that women are more susceptible to intense pain than men because of their lower pain tolerance. This is caused by biological factors and psychological factors that have a role in influencing differences in pain perception between the sexes [17].

According to the data in table 2, the majority of respondents (78.4%) reported only slight discomfort after the operation. Consistent with previous studies, this one describes the level of pain that patients felt following lower abdominal surgery at Sanglah General Hospital. Wiguna et al.

(2021) found that 71 out of 100 patients reported mild pain as their postoperative pain intensified. Increased pain can be influenced by the amount of surgery undergone. Postoperative pain appears due to mechanical stimulation of the wound which causes the body to produce chemical mediators of pain. The intensity of pain felt by patients varies greatly from mild pain to severe pain, this will decrease as recovery time progresses [5].

Table 3 displays the findings for sleep quality, which indicate that the majority of respondents (84.3% or 43 people) had satisfactory sleep. In a study by Sari et al. (2022), researchers looked at the correlation between postpartum pain and sleep quality in women who had cesarean sections. The majority of the participants (22 out of 33) reported good sleep quality, which is consistent with our findings. In some diseases, pain forces patients to fall asleep in unusual positions. In addition, there are several factors that can affect sleep such as disease, environment, stress, stimulants that cause someone to have trouble sleeping [18].

Based on the observations of researchers in several fracture cases, patients are required to use casts, which can affect pain intensity when performing early mobilization and interfere with comfort during sleep. Likewise, the environment and lighting factors that each individual has different comfort. Health status both physiological health conditions and psychological health greatly affects their sleep needs [19]. Sleep quality can be said to play a role in the post-surgical recovery process [20–23]. A good night's sleep is one in which a person does not wake feeling drowsy or otherwise disturbed. Difficulty maintaining a steady state of mind and body could result from poor quality sleep [24].

The data shown above indicates a substantial association (correlation) between the variable postoperative pain

intensity and sleep quality in patients following spinal anesthesia, with a correlation value of 0.756**. At the 0.01% level of significance, the association is indicated with an asterisk (**). The findings show a positive correlation value of 0.756, indicating a unidirectional association between the two variables. This suggests that a greater severity of postoperative pain is associated with lower sleep quality [25–27]. According to the results shown earlier, a significant relationship (mean) exists between the variable post-surgical pain intensity and sleep quality in patients following spinal anesthesia, as indicated by the Sig. (2-tailed) value of 0.000, which is less than 0.05 or 0.01.

This study confirms the findings of regarding the correlation between postoperative pain, anxiety, and sleep quality in the operating room. The results show that postoperative pain significantly affects sleep quality in patients at Budi Asih Hospital in Serang City in 2021 (p-value 0.001). Researchers assume that postoperative patients after 24 hours perceive pain from no pain to moderate pain. No pain to mild pain can be influenced by the type of surgery such as ROI (Remove of Implant), Up K-wires, nail extraction, vulvar biopsy and other minor surgeries, while moderate pain is influenced by the type of major surgery such as ORIF (Open reduction and internal fixation), hysterectomy, laparotomy, amputation and others [28–30].

The use of drugs as a way to reduce pain is the last alternative if the pain becomes increasingly severe, the patient can no longer bear the pain or the pain lasts for a long time [19]. Actual and prospective tissue damage causes patients to endure pain, which is an unpleasant sensory and emotional experience [5]. Postoperative pain can affect sleep quality, but in some people, pain does not affect sleep quality due to the different perceptions of each

patient and the level of need for sleep that varies with each individual [31]. Pain is one of the things that might disrupt a good night's sleep. Insomnia may be brought on by physical ailments [18].

Therefore, the role of medical personnel and the environment is very influential on patient comfort during the post-surgical recovery period in the hospital. Various efforts that can be made such as relaxation techniques, distraction, listening to music, and distractions that focus on feeling pain will help patients to relieve pain. During this time, pharmacological therapy is still given as a collaborative measure that can create a sense of comfort in post-surgical patients.

Limitations Research

Based on the researcher's experience in this research process, the limitations experienced are as follows: There is a combination in the delivery of the Numeric Rating Scale (NRS) with the delivery of the Verbal Rating Scale (VRS) where it is more dominant to use the NRS delivery, so that there are two different views. In the standard Pittsburgh Sleep Quality Index (PSQI) questionnaire there are differences in time, so it must be adopted and readjusted to the needs of the study. Researchers did not include the characteristics of the type of surgery, where these characteristics play a major role in the results of this study. Researchers did not look in detail regarding the pharmacological therapy that had been given according to the doctor's instructions, which could affect the intensity of the pain.

Conclusion

Understudy findings of RST Wijayakusuma Purwokerto, it could be concluded that most respondents who had spinal anesthesia after surgery experienced mild to moderate postoperative pain and reported good sleep quality in general.

Moreover, there was a significant relationship between postoperative pain intensity and sleep quality; it shows that those with lower pain levels experienced better sleep quality. This emphasizes the importance of good postoperative pain management in recovery, such as improving sleep quality in patients following spinal anesthesia.

Acknowledgment

The authors express sincere gratitude to Universitas Harapan Bangsa for its invaluable academic support throughout this research. We also extend our appreciation to Hospital RST Wijayakusuma Purwokerto for facilitating data collection and providing the necessary resources for this study.

Funding Information

None

Conflict of Interest Statement

The authors have confirmed that they have no competing interests.

Data Availability

The datasets used or generated in this study are available from the corresponding author upon reasonable request.

Author Contributions

Shelina: Conception and design of the study, Search Data Base, Methodology, Analysis Risk of Bias, Data Analysis and Interpretation, Writing, Review, and Editing. **Amin Susanto:** Study conception and design, search database, methodology, data analysis, and interpretation, and writing, review, and editing. **Septian Mixrova Sebayang** Conception and design of the study, Search Database, Methodology, Data Analysis, and Interpretation, Writing, Review, and Editing.

References

1. Budiyo A, Hamdiah D. Hubungan Antara Nyeri Dan Kecemasan Dengan Kualitas Tidur Pada Pasien Pasca Operasi Di Ruang Bedah. Ghidza J Gizi Dan Kesehat. 2022;6(2):191–9.
2. Saputri Gar, Nofita, Tiwi Ts. Rasionalitas Penggunaan Obat Anestesi Pada Sentral Rumah Sakit Pertamina Bintang Amin Lampung Tahun 2019. J Farm Malahayati. 2021;4(2):194–204.
3. Oroh A, Yudono Dt, Siwi As. Pengaruh Elevasi Kaki Terhadap Tekanan Darah Pada Pasien Sectio Caesaria Dengan Spinal Anestesi Di Instalasi Kamar Bedah Rumah Sakit Tk.Ii Robert Wolter Mongisidi Manado. J Inov Penelit. 2022;Vol.3 No.7(7):6857–64.
4. Hidayat S, Olfah Y, Sugeng. Pengaruh Terapi Distraksi Berdoa Terhadap Intensitas Nyeri Pada Pasien Pasca Bedah Anestesi Tulang Belakang Di Ruangan Pemulihan Rsud Andi Sulthan Daeng Kabupaten Bulukumba Sulawesi Selatan. Anaesth Nurs J. 2022;1(January 2021):42–8.
5. Mawaddah Ds. Hubungan Nyeri Terhadap Pola Tidur Pasien Post Operasi Appendisit Di Rsud Teungku Peukan Aceh Barat Daya. 2021;
6. Rohmayani Ms, Suwito J. Intensitas Nyeri Klien Pasca Operasi Mayor Di Rumah Sakit Islam Jemursari Surabaya. J Keperawatan. 2019;Xii(1):23–9.
7. Wiguna Kaa, Aribawa Ignm, Aryabiantara Iw, Senapathi Tga. Gambaran Intensitas Nyeri Pasien Pasca-Operasi Abdomen Bawah Di

- Rsup Sanglah. *J Med Udayana*. 2021;10(8):104–10.
8. Wainsani S, Khoiriyah K. Penurunan Intensitas Skala Nyeri Pasien Appendiks Post Appendiktomi Menggunakan Teknik Relaksasi Benson. *Ners Muda*. 2020;1(1):68–77.
 9. Baroki Saragih Mn, Lintang Suryani R, Burhan A. Overview Of Spinal Anesthesia Injection Techniques At Rsi Fatimah Cikacap: English. *Java Nurs J*. 2023 Oct 16;1(2):97–102.
 10. Fitriadi Y. Effort To Control Pulmonary Tuberculosis (Tb) In The Community Through Tuberculosis Alert Health Cadre Training. *J Community Empower Health*. 2023 Dec 8;6(3):133.
 11. Van Dijk Jfm, Zaslansky R, Van Boekel Rlm, Cheuk-Alam Jm, Baart Sj, Huygen Fjpm, Et Al. Postoperative Pain And Age: A Retrospective Cohort Association Study. *Anesthesiology*. 2021;135(6):1104–19.
 12. Badriah Dl, Negara Ap, Nur As. Faktor-Faktor Yang Berhubungan Dengan Kualitas Tidur Pada Klien Post Operasi Bedah Mayor Di Ruang Bedah Kelas Iii Rsud 45 Kuningan Tahun 2014. *J Kesehat Indra Husada*. 2017;5(2012):1–6.
 13. Maharani My, Masfuri M, Maria R. Analisis Faktor Yang Mempengaruhi Kualitas Tidur Pasien Ortopedi Pasca Pembedahan Yang Menjalani Rawat Inap. *J Pendidik Keperawatan Indones*. 2020;6(2):178–86.
 14. Nurdiansyah P, Susanto A, Burhan A, Suandika M, Wijayanti I. Overview Of The Incidence Of Post Operative Nausea And Vomiting In Spinal Anesthesia For Sectio Caesarea Patients In The Recovery Room Of Fatimah Cilacap Islamic Hospital. *Java Nurs J*. 2024 Feb 1;2(1):50–6.
 15. Susanti I, Subiyanto P, Burhan A. Overview Of Management Documentation Hypertension Patients: A Qualitative Study. 2023;16(03).
 16. Utami S. Efektivitas Aromaterapi Bitter Orange Terhadap Nyeri Post Partum Sectio Caesarea. *Unnes J Public Health*. 2016;5(4):316.
 17. Hidayati Hb, Amelia Egf, Turchan A, Rehatta Nm, Atika, Hamdan M. Pengaruh Usia Dan Jenis Kelamin Pada Skala Nyeri Pasien Trigeminal Neuralgia. *Aksona*. 2022;1(2):53–6.
 18. Noviyanti Ha, Sutrisna M, Kusmiran E. Hubungan Intensitas Nyeri Dengan Kualitas Tidur Pada Pasien Pasca Bedah Sesar. *J Persat Perawat Nas Indones*. 2019;4(2):59–66.
 19. Asdar F. Hubungan Intensitas Nyeri Dengan Kualitas Tidur Pada Pasien Post Operasi Laparatomi Di Rsud Labuang Baji Makassar. *J Ilm Kesehat Diagn*. 2018;12:509–13.
 20. Habsah Jumma Mh, Burhan A, Heri Susanti I. The Effectiveness Of Hyperbaric Bupivacaine Administration On Hemodynamic Changes In Sectio Caesarea Patients. *Java Nurs J*. 2024 Oct 1;2(3):247–56.
 21. Ayu Kartika Sari V, Suandika M, Burhan A, Tri Yudono D. Overview Of Induction Drug Dosage Types With The Eracs (Enhanced Recovery After Caesarean Surgery) Sectio Caesarean Method In Post Sectio Caesarean Patients At Hermina General. *Java Nurs J*. 2024 Jun 21;2(2):110–5.

22. Yanrin Asnp, Burhan A, Sukmaningtyas W, Adriani P. Gambaran Kejadian Hipotermi Pada Pasien Sectio Caesarea Di Rumah Sakit Umum Daerah Brebes. 2024 May 26 [Cited 2024 Aug 22]; Available From: <https://zenodo.org/doi/10.5281/zenodo.11313844>
23. Afrilya To, Sukmaningtyas W, Burhan A. Overview Of Post-Operative Nausea And Vomiting (Ponv) In Patients Undergoing Caesarean Section With Spinal Anesthesia At Fatimah Islamic Hospital Cilacap. 2023;16.
24. Annelia R, Sunarsih, Astuti T, Murhan A, Sulastrri. Pengaruh Terapi Akupresur Terhadap Kualitas Tidur Pasien Pos Operasi Sectio Caesarea Di Rsia Anugerah Medical Center Kota Metro. Mahesa Malahayati Health Stud J. 2022;2(3):611–21.
25. Susanti I, Mixrova Sebayang S, Burhan A. Impact Of Ondancentron In Intra-Anesthesia During Caesarean Section: A Meta-Analysis Of Randomized Trials. Java Nurs J. 2024 Feb 1;2(1):26–42.
26. Alverina F, Suryani Rl, Burhan A. An Overview Caring Behavior Of Anesthesiologist In Spinal Anesthesia Patients At Rsud Cilacap. Java Nurs J. 2024 Jun 21;2(2):124–30.
27. Vika Andriyani, Roro Lintang Suryani, Asmat Burhan. The Relationship Of Family Support With The Level Of Pre-Anesthesian Sectio Patients In The Hospital Pekalongan Regional General. Java Nurs J. 2024 Feb 1;2(1):43–9.
28. Triyadi F, Mixrova Sebayang S, Burhan A, Dwi Agus Yulianto, Refa Teja Muti. The Relationship Between Age And Duration Of Surgery With The Incidence Of Post Anesthesia Shivering In Section Caesarean Patients At Bendan Hospital Pekalongan City. Java Nurs J. 2024 Feb 1;2(1):7–14.
29. Permata Pp, Burhan A, Handayani Rn. Pengaruh Pemberian Aromaterapi Peppermint Terhadap Post Operative Nausea And Vomiting (Ponv) Post Operasi Spinal Anestesi Di Rsud 45 Kuningan. J Inov Glob. 2024 Oct 27;2(10):1517–34.
30. Fauzan Dh, Sebayang Sm, Burhan A, Suhendro. Gambaran Kejadian Shivering Post Spinal Anestesi Pada Pasien Benign Prostatic Hyperplasia Di Rumah Sakit Umum Daerah Cilacap. 2024 May 10 [Cited 2024 Aug 22]; Available From: <https://zenodo.org/doi/10.5281/zenodo.11173343>
31. Rokhima V, Sari Y. Hubungan Intensitas Nyeri Dengan Strategi Manajemen Nyeri Pada Pasien Fraktur Post Operasi Orif Di Rsu Setia Budi. J Vocat Health Sci. 2022;1(1):24–33.